GOVERNMENT OF NAGALAND DIRECTORATE OF HEALTH AND FAMILY WELFARE **NAGALAND: KOHIMA**

NO. DHFW/COVID-19/2019-20/2005 - OF Dated Kohima, the 25 th June 2020

To,

1. The Deputy Commissioner & Chairman, DTF of all districts

2. The Chief Medical Officer of all districts

3. The Medical Superintendent of all district hospitals / COVID Hospitals

Subject: Proper use of the recommended PPE by frontline workers in various settings

Sir/Madam.

In order to protect frontline workers from infection in the various workplace settings, a guideline was issued (Guidelines on rational use of PPE) where advice on the type of PPE to use in various settings (Point of Entry, Hospitals, Community Settings, Quarantine Centres, Home Quarantines) is given. The guideline is enclosed again for reference and implementation.

All frontline workers should use the appropriate PPE for the correct setting, and follow guideline on the correct way to use and dispose. Any breach of PPE should be immediately reported so that immediate segregation and testing can be done as per protocol.

Yours faithfully (DR. VIZOLIE Z.

Principal Director Directorate of Health and Family Welfare Nagaland : Kohima

NO. DHFW/COVID-19/2019-20/

Dated Kohima, theth June 2020

Copy to:

- 1. The Deputy Secretary to Chief Secretary, Government of Nagaland for information.
- 2. The Principal Secretary and Home Commissioner, for information.
- 3. The Commissioner and Secretary, Health and Family Welfare Department for information
- 4. Office Copy / Guard File

(DR. VIZOLIE Z. SUOKHRIE) **Principal Director** Directorate of Health and Family Welfare Nagaland : Kohima

Ministry of Health and Family Welfare Directorate General of Health Services [Emergency Medical Relief]

Novel Coronavirus Disease 2019 (COVID-19): Guidelines on rational use of Personal Protective Equipment

1. About this guideline

This guideline is for health care workers and others working in points of entries (POEs), quarantine centers, hospital, laboratory and primary health care / community settings. The guideline uses setting approach to guide on the type of personal protective equipment to be used in different settings.

2. Introduction

Coronaviruses are a large family of viruses, some causing illness in people and others that circulate among animals, including camels, cats and bats. Rarely, animal coronaviruses can evolve and infect people and then spread between people such as has been seen with MERS and SARS.

The outbreak of Novel coronavirus disease (now named COVID-19) was initially noticed from a seafood market in Wuhan city in Hubei Province of China in mid-December, 2019, has spread to more than 185 countries/territories worldwide including India.

The causative agent for COVID-19, earlier termed provisionally as novel Coronavirus has been officially named as SARS-CoV-2.

3. Mode of transmission

There is clear evidence of human-to-human transmission of SARS-CoV-2. It is thought to be transmitted mainly through respiratory droplets that get generated when people cough, sneeze, or exhale. SARS-CoV-2 also gets transmitted by touching, by direct touch and through contaminated surfaces or objects and then touching their own mouth, nose, or possibly their eyes. Healthcare associated infection by SARS-CoV-2 virus has been documented among healthcare workers in many countries.

The people most at risk of COVID-19 infection are those who are in close contact with a suspect/confirmed COVID-19 patient or who care for such patients.

4. **Personal Protective Equipment (PPE)**

Personal Protective Equipments (PPEs) are protective gears designed to safeguard the health of workers by minimizing the exposure to a biological agent.

4.1 Components of PPE

Components of PPE are goggles, face-shield, mask, gloves, coverall/gowns (with or without aprons), head cover and shoe cover. Each component and rationale for its use is given in the following paragraphs:

4.1.1 Face shield and goggles

Contamination of mucous membranes of the eyes, nose and mouth is likely in a scenario of droplets generated by cough, sneeze of an infected person or during aerosol generating procedures carried out in a clinical setting. Inadvertently touching the eyes/nose/mouth with a contaminated hand is another likely scenario. Hence protection of the mucous membranes of the eyes/nose/mouth by using face shields/ goggles is an integral part of standard and contact precautions. The flexible frame of goggles should provide good seal with the skin of the face, covering the eyes and the surrounding areas and even accommodating for prescription glasses.

4.1.2 Masks

Respiratory viruses that includes Coronaviruses target mainly the upper and lower respiratory tracts. Hence protecting the airway from the particulate matter generated by droplets / aerosols prevents human infection. Contamination of mucous membranes of the mouth and nose by infective droplets or through a contaminated hand also allows the virus to enter the host. Hence the droplet precautions/airborne precautions using masks are crucial while dealing with a suspect or confirmed case of COVID-19/performing aerosol generating procedures.

Masks are of different types. The type of mask to be used is related to particular risk profile of the category of personnel and his/her work. There are two types of masks which are recommended for various categories of personnel working in hospital or community settings, depending upon the work environment:

- 1. Triple layer medical mask
- 2. N-95 Respirator mask

4.1.2.1 Triple layer medical mask

A triple layer medical mask is a disposable mask, fluid-resistant, provide protection to the wearer from droplets of infectious material emitted during coughing/sneezing/talking.

4.1.2.2. N-95 Respirator mask

An N-95 respirator mask is a respiratory protective device with high filtration efficiency to airborne particles. To provide the requisite air seal to the wearer, such masks are designed to achieve a very close facial fit.

Such mask should have high fluid resistance, good breathability (preferably with an expiratory valve), clearly identifiable internal and external faces, duckbill/cup-shaped structured design that does not collapse against the mouth.

If correctly worn, the filtration capacity of these masks exceeds those of triple layer medical masks. Since these provide a much tighter air seal than triple layer medical masks, they are designed to protect the wearer from inhaling airborne particles.

4.1.3 Gloves

When a person touches an object/surface contaminated by COVID-19 infected person, and then touches his own eyes, nose, or mouth, he may get exposed to the virus. Although this is not thought

to be a predominant mode of transmission, care should be exercised while handling objects/surface potentially contaminated by suspect/confirmed cases of COVID-19.

Nitrile gloves are preferred over latex gloves because they resist chemicals, including certain disinfectants such as chlorine. There is a high rate of allergies to latex and contact allergic dermatitis among health workers. However, if nitrile gloves are not available, latex gloves can be used. Non-powdered gloves are preferred to powdered gloves.

4.1.4 Coverall/Gowns

Coverall/gowns are designed to protect torso of healthcare providers from exposure to virus. Although coveralls typically provide 360-degree protection because they are designed to cover the whole body, including back and lower legs and sometimes head and feet as well, the design of medical/isolation gowns do not provide continuous whole-body protection (e.g., possible openings in the back, coverage to the mid-calf only).

By using appropriate protective clothing, it is possible to create a barrier to eliminate or reduce contact and droplet exposure, both known to transmit COVID-19, thus protecting healthcare workers working in close proximity (within 1 meter) of suspect/confirmed COVID-19 cases or their secretions.

Coveralls and gowns are deemed equally acceptable as there is a lack of comparative evidence to show whether one is more effective than the other in reducing transmission to health workers. Gowns are considerably easier to put on and for removal. An apron can also be worn over the gown for the entire time the health worker is in the treatment area. Coveralls/gowns have stringent standards that extend from preventing exposure to biologically contaminated solid particles to protecting from chemical hazards.

4.1.5 Shoe covers

Shoe covers should be made up of impermeable fabric to be used over shoes to facilitate personal protection and decontamination.

4.1.6. Head covers

Coveralls usually cover the head. Those using gowns, should use a head cover that covers the head and neck while providing clinical care for patients. Hair and hair extensions should fit inside the head cover.

The specifications for all the PPEs are at Annexure-A.

5. Rational use of PPE

The PPEs are to be used based on the risk profile of the health care worker. The document describes the PPEs to be used in different settings.

5.1. Point of Entry

S. No.	Setting	Activity	Risk	Recommended PPE	Remarks
1	Health Desk	Provide information to travellers	Low risk	Triple layer medical mask Gloves	Minimum distance of one meter needs to be maintained.
2	Immigration counters, customs and airport security	Provide services to the passengers	Low risk	Triple layer medical mask Gloves	Minimum distance of one meter needs to be maintained.
3	Temperature recording station	Record Temperature with hand held thermal recorder.	Low risk	Triple layer medical mask Gloves	
4	Holding area/ Isolation facility of APHO/ PHO	Interview & Clinical examination by doctors/ nurses	Moderate Risk	N-95 masks Gloves	
5	Isolation facility of APHO	Clinical management (doctors, nurses)	Moderate Risk	N-95 masks Gloves	
		Attending to severely ill passenger	High risk	Full complement of PPE	When aerosol generating procedures are anticipated
5	Sanitary staff	Cleaning frequently touched surfaces/ Floor/ cleaning linen	Moderate risk	N-95 mask Gloves	
6	Administrative staff	Providing administrative support	No risk	No PPE	No contact with patients of COVID- 19. They should not venture into areas where suspect COVID-19 cases are being managed.

5.2. Hospital Setting

S. No	Setting	Activity	Risk	Recommended PPE	Remarks
1	Triage area	Triaging patients Provide triple layer mask to patient.	Moderate risk	N 95 mask Gloves	Patients get masked.
2	Screening area help desk/ Registration counter	Provide information to patients	Moderate risk	N-95 mask Gloves	
3	Temperature recording station	Record temperature with hand held thermal recorder	Moderate Risk	N 95 mask Gloves	
4	Holding area/ waiting area	Nurses / paramedic interacting with patients	Moderate Risk	N 95 mask Gloves	Minimum distance of one meter needs to be maintained.
5	Doctors chamber	Clinical management (doctors, nurses)	Moderate Risk	N 95 mask Gloves	No aerosol generating procedures should be allowed.
6	Sanitary staff	Cleaning frequently touched surfaces/ Floor/ cleaning linen	Moderate risk	N-95 mask Gloves	
7	Visitors accompanying young children and elderlies	Support in navigating various service areas	Low risk	Triple layer medical mask	No other visitors should be allowed to accompany patients in OPD settings. The visitors thus allowed should practice hand hygiene

5.2.1. Out Patient Department (Respiratory Clinic / Separate screening area)[#]

All hospitals should identify a separate triage and holding area for patients with Influenza like illness. If there is no triage area / holding area for patients due to resource constraints, such hospitals will follow the above guidance for general OPD.

5.2.2. In-patient Services

S. No.	Setting	Activity	Risk	Recommended PPE	Remarks
1	Individual isolation rooms/ cohorted isolation rooms	Clinical management	Moderate risk	N 95 mask Gloves	Patient masked. Patients stable. No aerosol generating activity.
2	ICU/ Critical	Critical care	High risk	Full complement of	Aerosol generating

	care	management		PPE	activities performed.
3	ICU /critical care	Dead body packing	High risk	Full complement of PPE	
4	ICU/ Critical care	Dead body transport to mortuary	Low Risk	Triple Layer medical mask Gloves	
5	Sanitation	Cleaning frequently	Moderate risk	N-95 mask	
		touched surfaces/ floor/ changing linen		Gloves	
6	Other Non- COVID treatment areas of hospital	Attending to infectious and non-infectious patients	Risk as per assessed profile of patients	PPE as per hospital infection prevention control practices.	No possibility of exposure to COVID patients. They should not venture into COVID-19 treatment areas.
7	Caretaker accompanying the admitted patient	Taking care of the admitted patient	Low risk	Triple layer medical mask	The caretaker thus allowed should practice hand hygiene, maintain a distance of 1 meter

5.2.3. Emergency Department

S.No	Setting	Activity	Risk	Recommended PPE	Remarks
1	Emergency	Attending emergency cases	Moderate risk	N 95 mask Gloves	When aerosol generating procedures are anticipated
2		Attending to severely ill patients of SARI	High risk	Full complement of PPE	Aerosol generating activities performed.

5.2.4. Pre-hospital (Ambulance) Services

S. No.	Setting	Activity	Risk	Recommended PPE	Remarks
1	Ambulance Transfer to designated hospital	Transporting patients not on any assisted ventilation	Moderate risk	N-95 mask Gloves	
		Management of SARI patient while transporting	High risk	Full complement of PPE	When aerosol generating procedures are anticipated
		Driving the ambulance	Low risk	Triple layer medical mask Gloves	Driver helps in shifting patients to the emergency

5.2.5. Other Supportive/ Ancillary Services

S. No.	Setting	Activity	Risk	Recommended PPE	Remarks
1.	Laboratory	Sample collection and transportation	High risk	Full complement of PPE	
		Sample testing	High risk	Full complement of PPE	
2	Mortuary	Dead body handling	Moderate Risk	N 95 mask Gloves	No aerosol generating procedures should be allowed. No embalming.
		While performing autopsy	High Risk	Full complement of PPE	No post-mortem unless until specified.
3	Sanitation	Cleaning frequently touched surfaces/ Floor/ cleaning linen in COVID treatment areas	Moderate risk	N-95 mask Gloves	
4	CSSD/Laundry	Handling linen of COVID patients	Moderate risk	N-95 mask Gloves	
5	Other supportive services	Administrative Financial Engineering Security, etc.	No risk	No PPE	No possibility of exposure to COVID patients. They should not venture into COVID-19 treatment areas.

5.3. Health Workers in Community Setting

S. No.	Setting	Activity	Risk	Recommended PPE	Remarks
1	ASHAs/ Anganwadi	Field Surveillance	Low Risk	Triple layer mask	Maintain distance of one meter.
	and other field staff			Gloves	Surveillance team to carry adequate triple layer masks to distribute to suspect cases detected on field surveillance
2	Doctors at supervisory level conducting field investigation	Field surveillance Clinical examination.	Medium risk	N 95 mask Gloves.	

5.4 Quarantine facility

S. No.	Setting	Activity	Risk	Recommended PPE	Remarks
1	Persons being quarantined		Low Risk	Triple layer mask	
2	Healthcare staff working at quarantine facility	Health monitoring and temperature recording	Low Risk	Triple layer mask Gloves	
		Clinical examination of symptomatic persons	Moderate Risk	N-95 masks Gloves	
3	Support staff		Low Risk	Triple layer mask Gloves	

5.5 Home Quarantine

S. No.	Setting	Activity	Risk	Recommended PPE	Remarks
1	Persons being quarantined		Low Risk	Triple layer mask	
2	Designated family member	Taking care of person being quarantined	Low Risk	Gloves	While cleaning commonly touched surfaces or handling soiled linen
3	Other family		No Risk	No PPE required	Maintain a distance of at least 1 meter from person under home quarantine. Senior citizens in the household should stay away from such persons under home quarantine.

Points to remember while using PPE

- 1. PPEs are not alternative to basic preventive public health measures such as hand hygiene, respiratory etiquettes which must be followed at all times.
- 2. Always (if possible) maintain a distance of at least 1 meter from contacts/suspect/confirmed COVID-19 cases
- 3. Always follow the laid down protocol for disposing off PPEs as detailed in infection prevention and control guideline available on website of MoHFW.

Annexure A

Personal Protection Equipment (PPE) - Specifications

(for Contact & Airborne precautions)

<u>1.</u> PPE Kit

1.1 Gloves

- Nitrile
- Non-sterile
- Powder free
- Outer gloves preferably reach mid-forearm (minimum 280 mm total length)
- Different sizes (6.5 & 7)
- Quality compliant with the below standards, or equivalent:
 - a. EU standard directive 93/42/EEC Class I, EN 455
 - b. EU standard directive 89/686/EEC Category III, EN 374
 - c. ANSI/SEA 105-2011
 - d. ASTM D6319-10
- **1.2** Coverall (medium and large)*
 - Impermeable to blood and body fluids
 - Single use
 - Avoid culturally unacceptable colors e.g. black
 - Light colors are preferable to better detect possible contamination
 - Thumb/fingerloops to anchor sleeves in place
 - Quality compliant with following standard
 - a. Meets or exceeds ISO 16603 class 3 exposure pressure, or equivalent
- **1.3** Goggles
 - With transparent glasses, zero power, well fitting, covered from all sides with elastic band/or adjustable holder.
 - Good seal with the skin of the face
 - Flexible frame to easily fit all face contours without too much pressure
 - Covers the eyes and the surrounding areas and accommodates for prescription glasses
 - Fog and scratch resistant
 - Adjustable band to secure firmly so as not to become loose during clinical activity
 - Indirect venting to reduce fogging
 - May be re-usable (provided appropriate arrangements for decontamination are in place) or disposable
 - Quality compliant with the below standards, or equivalent:
 - a. EU standard directive 86/686/EEC, EN 166/2002
 - b. ANSI/SEA Z87.1-2010

1.4. <u>N-95 Masks</u>

- Shape that will not collapse easily
- High filtration efficiency
- Good breathability, with expiratory valve
- Quality compliant with standards for medical N95 respirator:
 - a. NIOSH N95, EN 149 FFP2, or equivalent
- Fluid resistance: minimum 80 mmHg pressure based on ASTM F1862, ISO 22609, or equivalent
- Quality compliant with standards for particulate respirator that can be worn with full- face shield

1.5. Shoe Covers

- Made up of the same fabric as of coverall
- Should cover the entire shoe and reach above ankles
- 1.6. Face Shield
 - Made of clear plastic and provides good visibility to both the wearer and the patient
 - Adjustable band to attach firmly around the head and fit snuggly against the forehead
 - Fog resistant (preferable)
 - Completely covers the sides and length of the face
 - May be re-usable (made of material which can be cleaned and disinfected) or disposable
 - Quality compliant with the below standards, or equivalent:
 - a. EU standard directive 86/686/EEC, EN 166/2002
 - b. ANSI/SEA Z87.1-2010

3. Triple Layer Medical Mask

- Three layered medical mask of non-woven material with nose piece, having filter efficiency of 99% for 3 micron particle size.
 - a. ISI specifications or equivalent

4. Gloves

- Nitrile
- Non-sterile
- Powderfree
- Outer gloves preferably reach mid-forearm (minimum 280mm total length)
- Different sizes (6.5 & 7)
- Quality compliant with the below standards, or equivalent:
 - 1. EU standard directive 93/42/EEC Class I, EN 455
 - 2. EU standard directive 89/686/EEC Category III, EN 374
 - 3. ANSI/SEA 105-2011
 - 4. ASTM D6319-10

5. Body Bags - Specifications

- 1) Impermeable
- 2) Leak proof
- 3) Air sealed
- 4) Double sealed
- 5) Disposable
- 6) Opaque
- 7) White
- 8) U shape with Zip
- **9)** 4/6 grips
- **10)** Size: 2.2 x 1.2 Mts
- **11)** Standards:
 - a) ISO 16602:2007
 - b) ISO 16603:2004
 - c) IS016604:2004
 - d) ISO/DIS 22611:2003

All items to be supplied need to be accompanied with certificate of analysis from national/ international organizations/labs indicating conformity to standards

All items: Expiry 5 years

* Due to scarcity of coveralls, and risk versus benefit, that as an emergency temporary measure in larger public interest, in present given circumstances, the fabric that cleared/passed 'Synthetic Blood Penetration Resistance Test' (ISO 16603) and the garment that passed 'Resistance to penetration by biologically contaminated solid particles (ISO 22612:2005) may be considered as the benchmark specification to manufacture Coveralls." The Coveralls should be taped at the seams to prevent fluid/droplets/aerosol entry.

The test for these two standards (ISO 16603 and ISO 22612:2005), which can be performed in Indian laboratories are as per WHO Disease Commodity Package (Version 4.0)